

Supplementary Material

Atazanavir inhibits SARS-CoV-2 replication and pro-inflammatory cytokine production

Natalia Fintelman-Rodrigues^{1,9#}, Carolina Q. Sacramento^{1,9#}, Carlyle Ribeiro Lima^{9,#}, Franklin Souza da Silva^{2,9}, André C. Ferreira^{1,3,9}, Mayara Mattos^{1,9}, Caroline S. de Freitas^{1,9}, Vinicius Cardoso Soares¹, Suelen da Silva Gomes Dias¹, Jairo R. Temerozo^{4,5}, Milene D. Miranda⁶, Aline R. Matos⁶, Fernando A. Bozza^{7,8}, Nicolas Carels⁹, Carlos Roberto Alves², Marilda M. Siqueira⁶, Patrícia T. Bozza¹, Thiago Moreno L. Souza^{1,9,*}

- These authors contributed equally to this work

1 – Laboratório de Imunofarmacologia, Instituto Oswaldo Cruz (IOC), Fundação Oswaldo Cruz (Fiocruz), Rio de Janeiro, RJ, Brazil.

2 – Laboratório de Biologia Molecular e Doenças Endêmicas, IOC, Fiocruz, Rio de Janeiro, RJ, Brazil.

3 – Universidade Iguazu, Nova Iguaçu, RJ, Brazil.

4 – Laboratório de Pesquisas sobre o Timo, IOC, Fiocruz, Rio de Janeiro, RJ, Brazil.

5 - National Institute for Science and Technology on Neuroimmunomodulation (INCT/NIM), IOC, Fiocruz, Rio de Janeiro, RJ, Brazil.

6 – Laboratório de Vírus Respiratório e do Sarampo, IOC, Fiocruz, Rio de Janeiro, RJ, Brazil.

7 – Instituto Nacional de Infectologia Evandro Chagas, Fiocruz, Rio de Janeiro, RJ, Brazil

8 – Instituto D’or de Pesquisa e Ensino, Rio de Janeiro, RJ, Brazil

9 - National Institute for Science and Technology on Innovation in Diseases of Neglected Populations (INCT/IDPN), Center for Technological Development in Health (CDTS), Fiocruz, Rio de Janeiro, RJ, Brazil.

***Correspondence footnote:**

Thiago Moreno L. Souza, PhD

Fundação Oswaldo Cruz

Centro de Desenvolvimento Tecnológico em Saúde (CDTS)

Instituto Oswaldo Cruz (IOC)

Pavilhão Osório de Almeida, sala 16

Av. Brasil 4365, Manguinhos, Rio de Janeiro - RJ, Brasil, CEP 21060340

Tel.: +55 21 2562-1311

Email: tmoreno@cdts.fiocruz.br

Additional Results *in silico*

Molecular Dynamics

The enzyme-LPV complex was stable with a RMSD value not exceeding 3 Å, but when considering LPV alone, its RMSD exceeded 4 Å over the entire simulation (Figure S1A). By contrast, when complexed with ATV, the RMSD of SCV2-MP reached as much as 4 Å in the last 20 ns of the simulation, while the RMSD of ATV alone did not exceed 3 Å over the 100 ns (Figure S1B). After the step of energy minimization, at the beginning of the molecular dynamics, LPV presented 4 hydrogen bonds with the amino acids Thr25, Ser46, Gly143, and Gln189 (Figures S2A and S3A). After 100 ns of molecular dynamics, LPV interacted with Ser46 HIS41 (Figures 3A and 4A).

ATV's interaction, after energy minimization, occurred with residue Gln189 (Figure S2B and S3B), but after 100 ns of molecular dynamics, Glu166 acted as a hydrogen acceptor and His164 as a hydrogen donor (Figure 3B and 4B).

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68 Tables

69 **Table S1.** Interactions from Atazanavir (ATV) with the residues from the active site.

Docking Interactions of ATA		
Ligand	Residues	Distance, Å
ATV:H32	ASN142:OD1	1.91
ATV:H32	HIS164:O	2.64
ATV:H8	GLU166:OE1	2.44

70

Table S2. Interactions from Lopinavir (LPV) with the residues from the active site before (*) and after MD.

Molecular Dynamics Interactions of LPV		
Ligand	Residues	Distance, Å
LPV:O1	THR25:HG1	1.99*
LPV:O2	SER45:HG	1.75*
LPV:O	GLY143:H	2.31*
LPV:O2	GLN189:HE21	1.95*
LPV:O2	SER46:HD	1.70
LPV:O1	HIS41:HE2	2.03

Table S3. Interactions from Atazanavir (ATV) with residues from the active site before (*) and after MD.

Molecular Dynamics Interactions of ATV		
Ligand	Residues	Distance, Å
ATV:H28	GLN189:OE1	2.04*
ATV:H31	HIE167:O	1.87
ATV:H2	GLU166:O	2.51

Legend for the Figures

Figure S1. A representative structure of Mpro (PDB:6LU7) was color coded to show the electrostatic potential of residues in the active site for negative (blue) and positive (red) charges. Panel A, the cavities of ligand interaction designated S1*, S1 and S2 in the absence of inhibitors. Panel B, placement of LPV (cyan) docked in the S1* and S2 regions of the active site. Panel C, placement of ATV (orange) docked in the S1* and S1 regions of the active site.

Figure S2. Two-dimensional (2D) representations of the interactions of LPV (A) and ATV (B) in the Mpro active site based on a molecular docking analysis. Two hydrogen bonds are predicted between ATV and Mpro.

Figure S3. Root Mean Square Deviation (RMSD) along 100 ns of MD simulation of the Mpro (black) when complexed with A) LPV (red) and B) ATV (red). (A) LPV (red) is rather unstable while Mpro (black) remains stable. In (B), it is Mpro (black) that is destabilized while ATV (red) remains stable along the MD simulation.

Figure S4. Representative images of the Root Mean Square Deviation (RMSD) along 100 ns of molecular dynamics (MD) simulation. LPV (A) and ATV (B) are positioned in the Mpro active site at the beginning of 100 ns simulation.

Figure S5. Two-dimensional (2D) representation of the interactions of LPV (A) and ATV (B) into Mpro's active site at the beginning of 100 ns RMSD.

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124


125


126

127

128

129






134

135


136

137 Figure S3

A




B




138

A




B




140

141

A



B



- | | | | |
|---|---|---|--|
| <ul style="list-style-type: none">Charged (negative)Charged (positive)GlycineHydrophobicMetal | <ul style="list-style-type: none">PolarUnspecified residueWaterHydration siteHydration site (displaced) | <ul style="list-style-type: none">H-bondHalogen bondMetal coordinationPi-Pi stacking | <ul style="list-style-type: none">Pi-cationSalt bridgeSolvent exposure |
|---|---|---|--|